Additional Information

Tips For Observing and Photographing Dragonflies

Dragonflies can be loosely lumped into categories of "perchers" or "flyers". Perchers, such as some of the skimmers, have a usual or favorite perch which they return to repeatedly. Some flyers have a usual route, a beat which they fly back and forth in the same area. You can prefocus your camera or binoculars on that area to get a close look. Some have a large, long beat and are more difficult to predict, so some luck and persistence are required to find them perched. Most odonates are most active in the middle of the day, but some are more active early and late, roosting most of the day in the shade. Be prepared, research and study your subjects first to know the preferred habitats. Binoculars, especially close-focus binoculars are a great aid for observing all nature subjects.

Approach slowly, getting closer and snapping away as you go (with today's digital cameras, you can always delete later), a good distant shot is better than none and will be of some aid in identifying later.

Try to align planes of focus; for example, a side view is usually good, so move to the side for a clear side view of the dragonfly. A "3/4 view" can show face and eyes as well as the side. Some field marks are more important than others in differentiating one species from another. Get more than one angle if possible.

Sun at your back is always good for photographic detail but be ready to use flash to "fill" in the detail on the shadow side if the subject is back-lit or in any dark situation.

©2010 ATBI Group

Cannot be modified or reproduced except by written permission of ATBI

Updated July 2010

About ATBI

This brochure was compiled and developed by the All Taxa Biodiversity Inventory (ATBI) group. ATBI is an on-going project in Tennessee State Parks and State Natural Areas. The goal of ATBI is to document all species in Tennessee State Parks, monitor long-term trends in specific taxonomic groups, and promote education about and preservation of all biodiversity with increased science and citizen education efforts. To access the ATBI database, go to www.tn.gov/environment/parks/atbi.

For more information about ATBI, please contact David Hill, 7th Floor, L & C Tower, 401 Church St., Nashville, TN 37243. Email david.r.hill@tn.gov. Phone 615-253-2455.

Dragonfly References

Beaton, Giff. 2007. Dragonflies and Damselflies of Georgia and the Southeast. A Wormsloe Foundation Nature Book.

Dunkle, Sidney. 2000. Dragonflies through Binoculars: A Field Guide to Dragonflies of North America. Oxford University Press.

Nikula, Blair et al. 2002. Stokes Beginner's Guide to Dragonflies. Little, Brown & Company.

http://www.odonatacentral.org/. Dragonfly Society of the Americas. This site has checklists of dragonflies by county.

Collecting of plants or animals in state parks and natural areas is prohibited. Please help preserve Tennessee's natural heritage by leaving plants and animals for others to enjoy.

DUNBAR CAVE STATE NATURAL AREA

401 Old Dunbar Cave Rd Clarksville, TN 37043-5725 Phone: (931) 648-5526 www.tn.gov/environment/parks/DunbarCave/

TENNESSEE STATE PARKS DIVISION OF RESOURCE MANAGEMENT

All Taxa Biodiversity Inventory (ATBI)

Dragonflies and Damselflies of Dunbar Cave State Natural Area and Montgomery County



Photo © Richard Connors

Welcome to the wonderful world of dragonflies at beautiful Dunbar Cave State Natural Area! Of the approximate 155 species of odonates (dragonflies and damselflies) in Tennessee, 28 have been documented in this park or Montgomery County.

Above is a picture of a Widow Skimmer (*Libellula luctuosa*), male. A common "percher", widow skimmers are found near lakes, ponds and slow sections of streams. Both males and females have the solid black wing stripes next to the body. You can remember this one by: "the Widow wears black next to the body". The female has a dark brown abdomen with a bright yellow stripe down the sides.

Dragonfly and Damselfly Checklist of Dunbar Cave State Natural Area and Montgomery County

amselflies (Zygoptera)	☐ Widow Skimmer	Indian culture. In more recent times the cave was used as an important communit center and even as a country music ban
road-winged Damselflies	☐ Great Blue Skimmer	
Calopterygidae)	☐ Blue Dasher	
☐ Ebony Jewelwing	☐ Eastern Amberwing	
☐ Smoky Rubyspot	☐ Common Whitetail	
ond Damselflies (Coenagrionidae)	☐ Blue-faced Meadowhawk	
☐ Blue-fronted Dancer	☐ Carolina Saddlebags	
□ Variable Dancer	□ Black Saddlebags	
☐ Powdered Dancer	Habitat/History	
☐ Blue-tipped Dancer	Dunbar Cave State Natural Area is a 110- acre area in Montgomery County. Its sig- nificant feature is a well-explored scenic and historic cave, which above ground is	
☐ Dusky Dancer		
☐ Familiar Bluet		
☐ Stream Bluet	surrounded by an upland hardwood forest.	
☐ Skimming Bluet	A stream exits the cave and has been impounded to form a small lake that is inhabited by many fish, turtles, and other wild-life. The cave is the habitat for several rare animals including the federally endangered gray bat (<i>Myotis grisescens</i>) and the southern cavefish (<i>Typhlichthys subterraneus</i>). Cave salamanders, crickets and other animals as well as fungi and bacteria live near the front of the cave in the area known as	Good spots
☐ Orange Bluet		*Over the lake and in open areas, look for
☐ Slender Bluet		patrolling dragonflies
☐ Fragile Forktail		*All along the shore of Swan Lake in vegetation (for damselflies) and on prominent twigs for "percher" dragonflies
☐ Rambur's Forktail		
☐ Eastern Forktail		*Along woodland trail in sunny spots, look for perchers, especially in the spring
ragonflies (Anisoptera)		
ragonflies Darners (Aeshnidae)	the twilight zone.	1 / 1 / 1
☐ Shadow Darner	Dunbar is one of the most prominent caves in an area where caves and sinkholes are common. This eight mile long cave has	Notes
☐ Common Green Darner		
meralds (Corduliidae)	historical, natural, archaeological, and geo-	
☐ Prince Baskettail	logical significance. Dunbar Cave formed as the Red River cut through the Mississippian age limestone and lowered the local water table about two or three million years ago. This enabled the flow of water	
kimmers (Libellulidae)		
☐ Halloween Pennant		
☐ Eastern Pondhawk	through small cracks and solution joints	
□ Slaty Skimmer	from the sinkhole plain above it to the Red River drainage. The slightly acidic water dissolved the limestone along its route over	